arXiv:1612.02910v1 [math.RT] 9 Dec 2016

SYMMETRY CLASSES OF TENSORS AND SEMI-DIRECT PRODUCT OF FINITE ABELIAN GROUPS

KIJTI RODTES AND KUNLATHIDA CHIMLA

Abstract. In the study of symmetry classes of tensors, finding examples of symmetry classes of tensors that possess an o*-basis is of considerable interest. There are only few classes of groups that have been provided a necessary and sufficient condition for having such a basis. There is no general criterion for any finite groups yet. In this note, we provide a necessary and sufficient condition for the existence of o*-basis of symmetry classes of tensors associated with semi-direct product of some finite abelian groups and, consequently, their wreath product.

1. Introduction

Many branches of pure and applied mathematics: combinatorial theory, matrix theory, operator theory, group representation theory, differential geometry, partial differential equations, quantum mechanics and other areas motivate the study of symmetry classes of tensors [3]. In particular, finding examples of (higher) symmetry classes of tensors that possesses a special basis (o*-basis) raised by B.Y. Wang and M.P. Gong in [17] is of considerable interest.

Necessary and sufficient conditions for the existence of o*-basis of symmetry classes of tensors are provided in few classes of groups; for example, dihedral groups, [9], dicyclic groups, [4], semi-dihedral groups, [11], non abelian groups of order pq, [14], some subgroups of full symmetric groups and some types of p-groups in [8]. For any finite group, there is a necessary condition provided in [16].

It is well known that a finite group G can be decomposed as a semi-direct product $G = A \rtimes_\phi H$ of a group A by H if and only if the short exact sequence

$$\{e\} \longrightarrow A \longrightarrow G \longrightarrow H \longrightarrow \{e\}$$

splits. Note also that not every short exact sequence splits and then some extension groups of A by H can not always be written as a semi-direct product but it is a subgroup of some wreath product $A \wr_\Omega H$, for some H-set Ω (universal embedding theorem). Wreath product is defined via semi-direct product and it can be used to construct a lot of interesting groups. However, many well known groups can be written as a semi-direct product of finite abelian groups; for example, dihedral groups, semi-dihedral groups, non-abelian

MSC(2000): Primary 20C15; Secondary 15A69
Keywords: Symmetry classes of tensors, o*-basis, Semi-direct product, Wreath product.

groups of order pq and Z-groups (group in which its Sylow subgroups are all cyclic and this groups have applications in the classification of finite groups, [18]). In this article, we provide a necessary and sufficient condition for the existence of o*-basis of symmetry classes of tensors associated with semi-direct product of some finite abelian groups and, consequently, their wreath product.

2. Preliminary

Let V be an n-dimensional complex inner product space with an orthonormal basis $B = \{e_1, \ldots, e_n\}$. Then, the m-folds tensor space $V^{\otimes m} := V \otimes V \otimes \cdots \otimes V$ has $B^{\otimes m} = \{e_\alpha^{\otimes} \,|\, \alpha \in \Gamma_{m,n}\}$ as an orthonormal basis (with respect to the induced inner product), where $\Gamma_{m,n}$ is the set of all sequences $\alpha = (\alpha_1, \ldots, \alpha_m)$, with $1 \leq \alpha_i \leq n$ and

$$e_\alpha^{\otimes} = e_{\alpha_1} \otimes \cdots \otimes e_{\alpha_m}.$$

Let G be a subgroup of S_m. Define the action of G on $\Gamma_{m,n}$ by

$$\alpha\sigma = (\alpha_{\sigma^{-1}(1)}, \ldots, \alpha_{\sigma^{-1}(m)}).$$

The space $V^{\otimes m}$ is a left $\mathbb{C}G$-module with the action given by $\sigma e^{\otimes} = e^{\otimes}_{\alpha\sigma^{-1}}$ for each $\sigma \in G, \alpha \in \Gamma_{m,n}$, extended linearly. Under this action, the induced inner product on $V^{\otimes m}$ is G-invariant. Let $O(\alpha) = \{\alpha\sigma \,|\, \sigma \in G\}$ be the orbit of α. We write $\alpha \sim \beta$ if α and β belong to the same orbit in $\Gamma_{m,n}$. Let Δ be a system of distinct representatives of the orbits. We denote by G_α the stabilizer subgroup of α, i.e., $G_\alpha = \{\sigma \in G \,|\, \alpha\sigma = \alpha\}$.

For any $\sigma \in G$, define the operator $P_\sigma : V^{\otimes m} \longrightarrow V^{\otimes m}$ on the m-folds tensor space by

$$P_\sigma(v_1 \otimes \cdots \otimes v_m) = v_{\sigma^{-1}(1)} \otimes \cdots \otimes v_{\sigma^{-1}(m)}.$$

Let $\mathrm{Irr}(G)$ be the set of all irreducible characters of G. For each $\chi \in \mathrm{Irr}(G)$, the symmetry classes of tensors associated with G and χ is the image of the symmetry operator

$$T(G, \chi) = \frac{\chi(e)}{|G|} \sum_{\sigma \in G} \chi(\sigma) P_\sigma,$$

and it is denoted by $V_\chi(G)$. We say that the tensor $T(G, \chi)(v_1 \otimes \cdots \otimes v_m)$ is a decomposable symmetrized tensor, and we denote it by $v_1 * \cdots * v_m$. The dimension of $V_\chi(G)$ is given by

$$\dim(V_\chi(G)) = \frac{\chi(e)}{|G|} \sum_{\sigma \in G} \chi(\sigma) n^{c(\sigma)},$$

where $c(\sigma)$ is the number of cycles, including cycles of length one, in the disjoint cycle factorization of σ; see [13].

The inner product on V induces an G-invariant inner product on $V^{\otimes m}$ and hence on $V_\chi(G)$ which satisfies

$$\langle v_1 * \cdots * v_m, u_1 * \cdots * u_m \rangle = \frac{\chi(e)}{|G|} \sum_{\sigma \in G} \chi(\sigma) \prod_{i=1}^{m} \langle v_i, u_{\sigma(i)} \rangle = \frac{\chi(e)}{|G|} d_\chi^G(A),$$

where $A_{ij} = \langle v_i, u_j \rangle$ and d_χ^G is the generalized matrix function defined by $d_\chi^G(A) := \sum_{\sigma \in G} \chi(\sigma) \prod_{i=1}^m A_{i\sigma(i)}$. If v_i's and u_j's are orthonormal vectors, then the formula becomes

$$\langle e_\alpha^*, e_\beta^* \rangle = \begin{cases} 0 & \text{if } \alpha \nsim \beta, \\ \frac{\chi(1)}{|G|} \sum_{\sigma \in G_\beta} \chi(\sigma h^{-1}) & \text{if } \alpha = \beta h. \end{cases} \tag{2.1}$$

In particular, for $\sigma_1, \sigma_2 \in G$ and $\alpha \in \Gamma_{m,n}$, we have

$$\langle e^*_\alpha, e^*_{\alpha\sigma}\rangle = \frac{\chi(e)}{|G|}\sum_{h\in G_\alpha}\chi(\sigma h). \tag{2.2}$$

Since $\sigma e^*_\alpha = e^*_{\alpha\sigma^{-1}}$ for each σ [10, Lemma 1.3] and the induced inner product is G invariant,

$$\langle e^*_{\alpha\sigma_1}, e^*_{\alpha\sigma_2}\rangle = \langle e^*_\alpha, e^*_{\alpha\sigma_2\sigma_1^{-1}}\rangle, \tag{2.3}$$

for each σ_1, σ_2 in G and $\alpha \in \Gamma_{m,n}$. Moreover, by (2.2),

$$\| e^*_\alpha \|^2 = \frac{\chi(e)}{|G|}\sum_{h\in G_\alpha}\chi(h).$$

By setting

$$\overline{\Delta} = \{\alpha \in \Delta \,|\, \sum_{h\in G_\alpha}\chi(h) \neq 0\},$$

we have $e^*_\alpha \neq 0$ if and only if $\alpha \in \overline{\Delta}$.

For $\alpha \in \overline{\Delta}$, $V^*_\alpha := \langle e^*_{\alpha\sigma} \,|\, \sigma \in G\rangle$ is called the orbital subspace of $V_\chi(G)$. By (2.1), it follows that

$$V_\chi(G) = \bigoplus_{\alpha\in\overline{\Delta}} V^*_\alpha$$

is an orthogonal direct sum. In [6], it is proved that

$$\dim(V^*_\alpha) \;=\; \frac{\chi(e)}{|G_\alpha|}\sum_{h\in G_\alpha}\chi(h). \tag{2.4}$$

Thus, if χ is linear, then $|\chi(h)| \leq 1$, for each $h \in G_\alpha$ and thus $\dim V^*_\alpha \leq \chi(e) = 1$ and in this case, the set $\{e^*_\alpha \,|\, \alpha \in \overline{\Delta}\}$ is an orthogonal basis of $V_\chi(G)$. An orthogonal basis which consists of the decomposable symmetrized tensors e^*_α is called an *orthogonal $*$-basis* or *o*-basis* for short. If χ is not linear, it is possible that $V_\chi(G)$ has no o*-basis. More details and motivation for the study of symmetry classes of tensors can be found in [13].

To prove the main result, we also need some facts about vanishing sum of roots of unity. For a given natural number $m = p_1^{a_1}p_2^{a_2}\cdots p_t^{a_t}$, suppose that $\epsilon_1, \epsilon_2, ..., \epsilon_k \in \mathbb{C}$ are mth roots of unity. The main result of T.Y. Lam and K. H. Leung in [12], asserts that

$$\textit{if } k \notin \mathbb{N}_0\langle\{p_1, \ldots, p_t\}\rangle, \textit{ then } \textstyle\sum_{i=1}^k \epsilon_i \neq 0, \tag{2.5}$$

where $\mathbb{N}_0\langle\{p_1, \ldots, p_t\}\rangle = \{k_1p_1 + \cdots + k_tp_t \,|\, k_i \in \mathbb{N}_0, \text{ for each } 1 \leq i \leq t\}$ and $\mathbb{N}_0$ is the set of nonnegative integers.

3. Semi-direct product of finite abelian groups and Characters

Let A, H be finite groups with identity e_A, e_H, respectively. Let $\phi : H \longrightarrow \mathrm{Aut}(A)$ be a group homomorphism and we denote $\phi(h)$ by ϕ_h, for each $h \in H$. Let $G = A \rtimes_\phi H$. As a set, $G = \{(a, h) \,|\, a \in A, h \in H\}$. Operation on the group G is given by

$$(a_1, h_1)(a_2, h_2) = (a_1\phi_{h_1}(a_2), h_1h_2),$$

for each (a_1, h_1) and (a_2, h_2) in G. The identity of the group G is (e_A, e_H) and the inverse of $(a, h) \in G$ is $(\phi_{h^{-1}}(a^{-1}), h^{-1})$. Note that $\{(a, e_H) \,|\, a \in A\} \cong A$ is a normal subgroup of G

and $\{(e_A, h) \,|\, h \in H\} \cong H$ is a subgroup of G. It is also well known that G is a semi-direct product of A by H if and only if the short exact sequence $\{e\} \longrightarrow A \longrightarrow G \longrightarrow H \longrightarrow \{e\}$ splits. Moreover, by Schur-Zassenhaus theorem, if G is a finite group and N is a normal subgroup whose order is coprime to the order of the quotient G/N, then the short exact sequence

$$\{e\} \longrightarrow N \longrightarrow G \longrightarrow G/N \longrightarrow \{e\}$$

splits, i.e., G is a semi-direct product of N by G/N.

Let Ω be an H-set (with the action ϕ) and let K be the direct product $K = \prod_{\omega\in\Omega} A_\omega$ of copies of $A_\omega = A$ by the set Ω. The elements of K can be seen as sequences (a_ω) of elements of A indexed by Ω with componentwise multiplication. Then the action ϕ of H on Ω extends in a natural way to an action ϕ_Ω of H on the group K by $\phi_\Omega(h)(a_\omega) := (a_{h^{-1}\omega})$. Then the wreath product of A by H indexed by Ω is defined by $K \rtimes_{\phi_\Omega} H$ and it is denoted by

$$A \wr_\Omega H.$$

By the universal embedding theorem (known as the Krasner-Kaloujnine embedding theorem), if G is an extension of A by H, then there exists a subgroup of the wreath product $A \wr_\Omega H$ (for some Ω) which is isomorphic to G.

Now, suppose that A is an abelian groups. Then, the set of all irreducible characters (all are linear) $\mathrm{Irr}(A) := A^\vee = \mathrm{Hom}(A, \mathbb{C})$ is isomorphic to A. Thus, the action of H on A induces the action on $A^\vee$. Precisely, for each $x \in A^\vee$ and $h \in H$, $h\cdot x := \phi_h(x) = x\circ\phi_h$. Let $[x]$ be the orbital and H_x be the stabilizer of x under this action. Let U be an irreducible representation of H_x. We define a representation $V_{([x],U)}$ of $G = A \rtimes_\phi H$ as follows. As a representation of H, we set

$$V_{([x],U)} := \mathrm{Ind}_{H_x}^H(U) = \{f : H \longrightarrow U \,|\, f(\sigma h) = \sigma f(h), \sigma \in H_x, h \in H\}.$$

Next, we introduce an additional action of A on $V_{([x],U)}$ by $(af)(h) := (x \circ \phi_h)(a)f(h)$. The combination of these two actions becomes an action of $G = A \rtimes_\phi H$.

It can be shown that (see Theorem 4.75, [5], for example) this representation is independent from the choice of $x \in [x]$. Also, if $\{[x_1], \ldots, [x_k]\}$ is the set of all disjoint orbits for the action on H on $A^\vee$, then

$$\{V_{([x_i],U)} \,|\, U \in \mathrm{Irr}(H_{x_i}), i = 1, 2, ..., k\},$$

forms a complete set of irreducible representations of $G = A \rtimes_\phi H$. The character χ_V of $V := V_{([x],U)}$, is given by the Mackey-type formula: for each $a \in A$ and $g \in H$,

$$\chi_V(a, g) = \frac{1}{|H_x|} \sum_{h\in H : hgh^{-1}\in H_x} x(\phi_h(a))\chi_U(hgh^{-1}). \tag{3.1}$$

In particular, if both A and H are abelian, then formula (3.1) reduces to

$$\chi_V(a, g) = \begin{cases} \frac{\chi_U(g)}{|H_x|} \sum_{h\in H} x(\phi_h(a)) & \text{if } g \in H_x, \\ 0 & \text{if } g \notin H_x. \end{cases} \tag{3.2}$$

4. Results

Let $G = A \rtimes_\phi H$, where both A, H are finite abelian groups, and χ be an irreducible character of G. Let V be an inner product space with $\dim V = n > 1$ and $V_\chi(G)$ be a symmetry class of tensors in the m-folds tensor space $V^{\otimes m}$. In the the following results, we denote by $\mathrm{Prime}(|A|)$ the set of all prime factors of $|A|$.

Theorem 4.1. *If there is $\alpha \in \Gamma_{m,n}$ such that $G_\alpha = \{e\}$ and $|H| \notin \mathbb{N}_0\langle \mathrm{Prime}(|A|)\rangle$, then $V_\chi(G)$ admits o*-basis if and only if χ is linear.*

Proof. We write χ for $\chi_{V_{([x],U)}}$, where $U \in \mathrm{Irr}(H_x)$. If χ is linear then, by discussion in Section 2, $V_\chi(G)$ admits o*-basis. On the other hand, assume that there is an o*-basis for $V_\chi(G)$. Then, $V^*_\alpha(G)$ has an o*-basis; say, $B = \{e^*_{\alpha(a_1,h_1)}, \ldots, e^*_{\alpha(a_{s_\alpha},h_{s_\alpha})}\}$ for some $(a_i, h_i) \in G$, where $s_\alpha = \dim V^*_\alpha(G)$.

Since H is abelian, U is linear. Then, by (3.2), $G_\alpha = \{e := (e_A, e_H)\}$, and by Freese's theorem, we calculate that

$$s_\alpha = \frac{\chi(e_A, e_H)}{|G_\alpha|} \sum_{(a,h)\in G_\alpha} \chi(a,h) = \left(\frac{\chi_U(e_H)}{|H_x|} \sum_{h\in H} x(\phi_h(e_A)) \right)^2 = (\frac{|H|}{|H_x|})^2. \tag{4.1}$$

Moreover, in (3.2), $\sum_{h\in H} x(\phi_h(a))$ is an $|H|$-terms sum of $|A|$th-roots of unity. Since $|H| \notin \mathbb{N}_0\langle \mathrm{Prime}(|A|)\rangle$, by (2.5) and (3.2), the sum can not be zero. We now conclude that, for each $(a, h) \in G$,

$$\chi(a,h) = 0 \text{ if and only if } h \notin H_x. \tag{4.2}$$

We claim that $h_i H_x \cap h_j H_x = \emptyset$ if $i \neq j$. Suppose $h_i = \sigma h_j$, for some $\sigma \in H_x$. Since $e^*_{\alpha(a_i,h_i)}$ and $e^*_{\alpha(a_j,h_j)}$ are orthogonal and the inner product is G invariant,

$$\begin{aligned}
0 &= \left\langle e^*_{\alpha(a_i,h_i)}, e^*_{\alpha(a_j,h_j)} \right\rangle \\
&= \left\langle e^*_{\alpha}, e^*_{\alpha(a_j,h_j)(a_i,\sigma h_j)^{-1}} \right\rangle \\
&= \left\langle e^*_{\alpha}, e^*_{\alpha(a_j,h_j)(\phi_{(\sigma h_j)^{-1}}(a_i^{-1}),(\sigma h_j)^{-1})} \right\rangle \\
&= \left\langle e^*_{\alpha}, e^*_{\alpha(a_j\phi_{\sigma^{-1}}(a_i^{-1}),\sigma^{-1})} \right\rangle \\
&= \frac{\chi(e)}{|G|} \chi(a_j\phi_{\sigma^{-1}}(a_i^{-1}), \sigma^{-1}) \neq 0,
\end{aligned}$$

by (4.2), which is a contradiction and then the claim becomes true. Therefore,

$$|H| \geq s_\alpha |H_x|.$$

By (4.1), we conclude that $|H_x| = |H|$. This implies that

$$\deg(\chi) = \chi(e_A, e_H) = |H|/|H_x| = 1$$

and hence χ is linear. □

Let Ω be a finite H-set. Then $\mathrm{Prime}(|A|^{|\Omega|}) = \mathrm{Prime}(|A|)$. Also, if A is abelian, then so is $K = \prod_{\omega\in\Omega} A_\omega$, where $A_\omega = A$, for each $\omega \in \Omega$. Hence, the following result is an immediate fact from Theorem 4.1.

Corollary 4.2. *Let $G = A \wr_\Omega H$, where A, H are finite abelian groups. If there is $\alpha \in \Gamma_{m,n}$ such that $G_\alpha = \{e\}$ and $|H| \notin \mathbb{N}_0\langle \text{Prime}(|A|)\rangle$, then $V_\chi(G)$ admits o*-basis if and only if χ is linear.*

Condition for having an o*-basis of symmetry classes of tensors associated to the following type of groups can be obtained from the above theorem as well.

Corollary 4.3. *Let V be an inner product space with $\dim V = n > 1$. The symmetry classe of tensors $V_\chi(G)$, associated to the groups G below, admits o*-basis if and only if χ is linear.*

(1) *Dihedral group D_{2s}, where s is odd.*
(2) *Non-abelian groups of order pq.*
(3) *Z-group G that its order has two prime factors and $G_\alpha = \{e\}$ for some $\alpha \in \Gamma_{m,n}$.*

Proof. (1) The dihedral group D_{2s} is a semi-direct product of $C_s = \langle a\rangle$ by $C_2 = \langle b\rangle$, i.e., $D_{2s} = C_s \rtimes_\phi C_2$, where $\phi : C_2 \longrightarrow \text{Aut}(C_s)$ is defined by $\phi_b(a) = a^{-1}$. Note that $|C_2| = 2 \notin \mathbb{N}_0\langle \text{Prime}(|C_s|)\rangle$ (because s is odd), and $G_\alpha = \{e\}$, where $\alpha = (1, 2, 1, ...1) \in \Gamma_{s,n}$ (cf. [9]). By Theorem 4.1, the conclusion is immediate.

(2) Each non-abelian group G of order pq, where q is prime and p divides $q-1$, is a semi-direct product of a cyclic group $C_q = \langle a\rangle$ by a cyclic group $C_p = \langle b\rangle$, i.e., $G = C_q \rtimes_\phi C_p$ such that $\phi : C_q \longrightarrow \text{Aut}(C_q)$ is given by $\phi_b(a) = a^r$, where r is a primitive root of the congruence $z^p \equiv 1 (\text{mod}\, q)$, [1]. It is clear that $|C_p| = p \notin \mathbb{N}_0\langle \text{Prime}(|C_q|)\rangle$ and $G_\alpha = \{e\}$, where $\alpha = (1, 2, 1, \ldots, 1)) \in \Gamma_{q,n}$, [14]. Now, the result is immediate by Theorem 4.1.

(3) Note that Z-group G is an extension group of a split short exact sequence of a cyclic group C_s by a cyclic group C_t, whose orders are relatively primes, [18]. So, it is a semi-direct product $G = C_s \rtimes_\phi C_t$, for some $\phi \in \text{Hom}(C_t, \text{Aut}(C_s))$. By the assumptions, all requirements of the Theorem 4.1 are satisfied and thus the result follows. □

In the following results, let G be any finite group. Let χ be an irreducible character of G, $V_\chi(G) \leq V^{\otimes m}$, and denote $Z_\chi := \{g \in G \,|\, \chi(g) = 0\}$. Using the same idea as in the proof of Theorem 4.1, we have:

Theorem 4.4. *Suppose that there is $\alpha \in \Gamma_{m,n}$ such that $G_\alpha = \{e\}$. If there exists a subgroup H of G such that $H \subseteq G \setminus Z_\chi$ and $[G : H] < \chi(e)^2$, then $V_\chi(G)$ does not admit an o*-basis.*

Proof. Since $G_\alpha = \{e\}$, $s_\alpha = \dim(V^*_\alpha(G)) = \chi(e)^2$. Suppose that $B = \{e^*_{\alpha g_1}, \ldots, e^*_{\alpha g_{s_\alpha}}\}$ is an o*-basis of $V^*_\alpha(G)$. We claim that $Hg_i \cap Hg_j = \emptyset$ if $i \neq j$. Suppose for a contradiction that $h_1 g_i = h_2 g_j$, for some $1 \leq i \neq j \leq s_\alpha$ and some $h_1, h_2 \in H$. Then, $g_j g_i^{-1} = h_2^{-1} h_1$ and thus

$$0 = \left\langle e^*_{\alpha g_i}, e^*_{\alpha g_j} \right\rangle = \left\langle e^*_\alpha, e^*_{\alpha g_j g_i^{-1}} \right\rangle = \left\langle e^*_\alpha, e^*_{\alpha h_2^{-1} h_1} \right\rangle = \frac{\chi(e)}{|G|} \chi(h_2^{-1} h_1) \neq 0,$$

by (2.2), (2.3) and $h_2^{-1} h_1 \in H \subseteq G \setminus Z_\chi$. Hence, $|G| \geq s_\alpha |H|$, which completes the proof. □

Acknowledgements

Most parts of this paper were done while the first author was the visiting assistant professor in the Department of Mathematics and Statistics, Auburn University, AL, USA. He would like to express his thanks to Prof. Tin-Yau Tam for the hospitality and his suggestion on the earlier draft.

References

[1] S. K. Berberian, *Non-abelian groups of order pq*, Amer. Math. Monthly 60 (1953), pp. 37-40.
[2] C. Bessenrodt, M. R. Pournaki, and A. Reifegerste, *A note on the orthogonal basis of a certain full symmetry class of tensors*, Linear Algebra Appl. 370 (2003), pp. 369–374.
[3] C. K. Li and A. Zaharia, *Induced operators on symmetry classes of tensors*, Trans. Amer. Math. Soc. 354 (2002), pp. 807-836.
[4] M. R. Darafsheh and M. R. Pournaki, *On the orthogonal basis of the symmetry classes of tensors associated with the dicyclic group*, Linear and Multilinear Algebra 47 (2000), pp. 137–149.
[5] P. Etingof, O. Golberg, S. Hensel. T. Liu, A. Schwendner, D. Vaintrob and E. Yudovina, *Introduction to representation theory* (MIT Open Courseware, 2011), http://math.mit.edu/ etingof/replect.pdf
[6] R. Freese, *Inequalities for generalized matrix functions based on arbitrary characters*, Linear Algebra Appl. 7 (1973), pp. 337–345.
[7] R. R. Holmes, *Orthogonal bases of symmetrized tensor spaces*, Linear and Multilinear Algebra 39 (1995), pp. 241–243.
[8] R. R. Holmes, *Orthogonality of cosets relative to irreducible character of finite groups*, Linear and Multilinear Algebra 52, (2004), pp. 133-143.
[9] R. R. Holmes and T. Y. Tam, *Symmetry classes of tensors associated with certain groups*, Linear and Multilinear Algebra 32 (1992), pp. 21–31.
[10] R. R. Holmes and A. Kodithuwakku, *Orthogonal bases of Brauer symmetry classes of tensors for the dihedral group*, Linear and Multilinear Algebra 61 (2013), pp. 1136–1147.
[11] M. Hormozi and K. Rodtes, *Symmetry classes of tensors associated with the semi-dihedral groups* SD_{8n}, Colloquium Mathematicum 131 (2013), 59–67.
[12] T. Y. Lam and K. H. Leung, *On vanishing sums of roots of unity*, J. Algebra 224 (2000), pp. 91-109.
[13] R. Merris, *Multilinear Algebra*, Gordan and Breach Science Publishers, Amsterdam, 1997.
[14] K. Rodtes, *Symmetry classes of tensors associated to nonabelian groups of order pq*, Bull. Aust. Math. Soc. 94 (2016), pp. 36-42.
[15] M. A. Shahabi, K. Azizi and M. H. Jafari, *On the orthogonal basis of symmetry classes of tensors*, J. Algebra 237 (2001), pp. 637–646.
[16] M. Shahryari, *On the orthogonal bases of symmetry classes*, J. Algebra 220 (1999), pp. 327-332.
[17] B. Y. Wang and M. P. Gong, *A higher symmetry class of tensors with an orthogonal basis of decomposable symmetrized tensors*, Linear and Multilinear Algebra 30 (1991), pp. 61-64.
[18] W. J. Wong, *On finite groups with semi-dihedral Sylow 2 subgroups*, J. Algebra 4 (1966), pp. 52-63.

Kijti Rodtes

Department of Mathematics, Faculty of Science,
Naresuan University, and Research Center for Academic Excellent in Mathematics
Phitsanulok 65000, Thailand
kijtir@nu.ac.th, and

Department of Mathematics and Statistics,
Auburn University, Alabama 36849, USA
kzr0033@auburn.edu

Kunlathida Chimla

Department of Mathematics, Faculty of Science,
Naresuan University, Phitsanulok 65000, Thailand
kunlathida_ nu@hotmail.com